

COMMON BOUNDARY REGULAR FIXED POINTS FOR HOLOMORPHIC SEMIGROUPS IN STRONGLY CONVEX DOMAINS

MARCO ABATE AND FILIPPO BRACCI[†]

Dedicated to Prof. David Shoikhet for his 60th anniversary

ABSTRACT. Let D be a bounded strongly convex domain with smooth boundary in $\mathbb{C}^N$. Let (ϕ_t) be a continuous semigroup of holomorphic self-maps of D . We prove that if $p \in \partial D$ is an isolated boundary regular fixed point for ϕ_{t_0} for some $t_0 > 0$, then p is a boundary regular fixed point for ϕ_t for all $t \geq 0$. Along the way we also study backward iteration sequences for elliptic holomorphic self-maps of D .

1. INTRODUCTION

Let $D \subset \mathbb{C}^N$ be a bounded domain. A continuous one-parameter semigroup of holomorphic self-maps of D (or, shortly a semigroup of holomorphic self-maps of D) is a family $(\phi_t)_{t \geq 0}$ of holomorphic self-maps of D such that $\phi_{t+s} = \phi_t \circ \phi_s$ for all $s, t \geq 0$, $\phi_0 = \text{id}_D$ and $\mathbb{R}^+ \ni t \mapsto \phi_t(z)$ is locally absolutely continuous locally uniformly in z . Namely, (ϕ_t) is a continuous semigroup morphism between the semigroup $(\mathbb{R}^+, +)$ endowed with the Euclidean topology and the composition semigroup of holomorphic self-maps of D endowed with the topology of uniform convergence on compacta. Every semigroup of holomorphic self-maps of D is generated by a $\mathbb{R}$ -semi-complete holomorphic vector field on D , called the infinitesimal generator associated with the semigroup.

Semigroups of holomorphic self-maps have been extensively studied (see, e.g. [27]), in connection with various areas of analysis, including geometric function theory, operator theory, iteration theory, theory of branching stochastic processes, Loewner theory.

The aim of this paper is to give a contribution to boundary dynamics of semigroups on a bounded strongly convex domain with smooth boundary $D \subset \mathbb{C}^n$, studying common boundary (regular) fixed points of semigroups in D (among which, a particular interesting case is the unit ball of $\mathbb{C}^N$).

As holomorphic self-maps of a domain might not extend continuously to the boundary, if $p \in \partial D$ and $f: D \rightarrow D$ is holomorphic, one might think of p as a boundary fixed point of f if f admits limit p along all sequences converging to p in some “admissible” subset of D (see Section 2 for precise definitions). In case $D = \mathbb{D}$ the unit disc of $\mathbb{C}$, such admissible subsets are exactly the Stolz angles, and thus $p \in \partial \mathbb{D}$ is a boundary fixed point of f if f has non-tangential limit p at p .

Boundary fixed points can be divided into two categories, the *boundary regular fixed points*—BRFP’s for short—and the *irregular* (or super-repulsive) fixed points.

[†]Supported by the ERC grant “HEVO - Holomorphic Evolution Equations” n. 277691.

In the unit disc, the first category is formed by those boundary fixed points for which the angular derivative of the map exists finitely. By the classical Julia-Wolff-Carathéodory theorem, the existence of the (finite) angular derivative at a boundary point for a holomorphic self-map of $\mathbb{D}$ corresponds to the finiteness of the so-called *boundary dilatation coefficient*—which, roughly speaking, measure the rate of approach of $f(z)$ to p as $z \rightarrow p$. In higher dimension, a boundary regular fixed point for a holomorphic self-map f of D is a point $p \in \partial D$ for which the admissible limit of f at p is p and the boundary dilation coefficient of f at p is finite.

Common boundary regular fixed points for a semigroup (i.e., boundary points which are BRFP's for each element of the semigroup) have been studied and characterized in terms of the local behavior of the associated infinitesimal generator in [13, 19, 15, 17, 5, 9].

Moreover, it is known (see, [12, Theorem 1], [13, Theorem 2], [28, pag. 255], [16]) that given a semigroup (ϕ_t) of holomorphic self-maps of $\mathbb{D}$, a point $p \in \partial \mathbb{D}$ is a boundary (regular) fixed point of ϕ_{t_0} for some $t_0 > 0$ if and only if it is a boundary (regular) fixed point of ϕ_t for all $t \geq 0$. The proof of this fact relies on the existence of the so-called Königs function, a univalent map from $\mathbb{D}$ to $\mathbb{C}$ which (simultaneously) linearizes the semigroup (ϕ_t) . Such a tool is not available in higher dimension in general, due essentially to the lack of a Riemann uniformization theorem.

The aim of this paper is to extend part of the previous results to higher dimension using an argument based on complex geodesics and backward iteration sequences. In order to state our result, we say that if $f: D \rightarrow D$ is holomorphic and $p \in \partial D$ is a BRFP with boundary dilatation coefficient $A > 0$, then p is *isolated* if there exists a neighborhood U of p such that $U \cap \partial D$ contains no BRFP's for f with boundary dilatation coefficients $\leq A$ except p .

Then our main result (proved in Section 6) can be stated as follows

Theorem 1.1. *Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary. Let (ϕ_t) be a semigroup of holomorphic self-maps of D . Suppose $p \in \partial D$ is an isolated boundary regular fixed point for ϕ_{t_0} for some $t_0 > 0$. Then p is a boundary regular fixed point for ϕ_t for all $t \geq 0$.*

The proof of Theorem 1.1 relies on the study of “regular contact points” for semigroups and backward iteration sequences. In particular, in Section 5, using complex geodesics, we prove that given a regular contact point $p \in \partial D$ for ϕ_{t_0} for some $t_0 > 0$, then p is a regular contact point for ϕ_t for all $t \in [0, t_0]$ and the curve $[0, t_0] \ni t \mapsto \phi_t(p) \in \partial D$ is continuous, extending to higher dimension one of the results in [11]. Along the way, in Section 3 we extend the results about existence and convergence of backward iteration sequences for holomorphic self-maps of D obtained in [3] to the case of “rotational elliptic maps”.

2. BRFP'S IN STRONGLY CONVEX DOMAINS

Let D be a bounded strongly convex domain in $\mathbb{C}^n$ with smooth boundary. A *complex geodesic* is a holomorphic map $\varphi: \mathbb{D} \rightarrow D$ which is an isometry between the Poincaré metric $k_{\mathbb{D}}$ of $\mathbb{D} = \{\zeta \in \mathbb{C} \mid |\zeta| < 1\}$ and the Kobayashi distance k_D in D ([20]). A holomorphic map $h: \mathbb{D} \rightarrow D$ is a complex geodesic if and only if it is an infinitesimal isometry between the Poincaré metric $\kappa_{\mathbb{D}}$ of $\mathbb{D}$ and the Kobayashi metric κ_D of D (see [1, Ch. 2.6]).

According to Lempert (see [21, 22, 23] and [1]), any complex geodesic extends smoothly to the boundary of the disc and $\varphi(\partial\mathbb{D}) \subset \partial D$. Moreover, given any two points $z, w \in \overline{D}$, $z \neq w$, there exists a complex geodesic $\varphi: \mathbb{D} \rightarrow D$ such that $z, w \in \varphi(\overline{\mathbb{D}})$. Such a geodesic is unique up to pre-composition with automorphisms of $\mathbb{D}$. Conversely, if $\varphi: \mathbb{D} \rightarrow D$ is a holomorphic map such that $k_D(\varphi(\zeta_1), \varphi(\zeta_2)) = k_{\mathbb{D}}(\zeta_1, \zeta_2)$ for some $\zeta_1 \neq \zeta_2 \in \mathbb{D}$, then φ is a complex geodesic.

Similarly, given $z \in D$ and $v \in T_z D \setminus \{0\}$, there exists a unique complex geodesic such that $\varphi(0) = z$ and $\varphi'(0) = \lambda v$ for some $\lambda > 0$.

If $\varphi: \mathbb{D} \rightarrow D$ is a complex geodesic then there exists a (unique when suitably normalized) holomorphic map $\tilde{\rho}: D \rightarrow \mathbb{D}$, smooth up to ∂D such that $\tilde{\rho} \circ \varphi = \text{id}_{\mathbb{D}}$. The map $\tilde{\rho}$ is called the *left inverse* of φ . It is known that $\tilde{\rho}^{-1}(e^{i\theta}) = \{\varphi(e^{i\theta})\}$ for all $\theta \in \mathbb{R}$, while the fibers $\tilde{\rho}^{-1}(\zeta)$ are the intersection of D with affine complex hyperplanes for all $\zeta \in \mathbb{D}$ (see, e.g., [10, Section 3]).

In the sequel we shall use the following result (see [10, Corollary 2.3, Lemma 3.5])

Proposition 2.1. *Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary and let $z_0 \in D$. Let $\{\varphi_k\}_{k \in \mathbb{N}}$ be a family of complex geodesics of D such that $\varphi_k(0) = z_0$ for all $k \in \mathbb{N}$ and let $\tilde{\rho}_k$ denote the left inverse of φ_k for $k \in \mathbb{N}$. If $\{\varphi_k\}$ converges uniformly on compacta of $\mathbb{D}$ to a function $\varphi: \mathbb{D} \rightarrow \mathbb{C}^N$, then φ is a complex geodesic and $\varphi_k \rightarrow \varphi$ uniformly in $\overline{\mathbb{D}}$. Moreover, $\{\tilde{\rho}_k\}$ converges uniformly in $\overline{D}$ to the left inverse $\tilde{\rho}$ of φ .*

Given $z_0 \in D$ and $p \in \partial D$, we will denote by $\varphi_p: \mathbb{D} \rightarrow D$ the unique complex geodesic such that $\varphi_p(0) = z_0$ and $\varphi_p(1) = p$ and by $\tilde{\rho}_p$ its left inverse. We will also denote

$$\rho_p := \varphi_p \circ \tilde{\rho}_p: D \rightarrow \varphi(\mathbb{D}).$$

We recall now the notion of “admissible limits” in strongly convex domain (see [1, 2]).

Definition 2.2. Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary and let $z_0 \in D$ and $p \in \partial D$. A sequence $\{z_k\} \subset D$ converging to p is said to be *special* if

$$\lim_{k \rightarrow \infty} k_D(z_k, \rho_p(z_k)) = 0.$$

The sequence $\{z_k\}$ is called *restricted* if $\tilde{\rho}_p(z_k) \rightarrow 1$ non-tangentially in $\mathbb{D}$. A continuous curve $\gamma: [0, 1] \rightarrow D$ such that $\gamma(1) = p$ is called *special*, respectively *restricted*, provided for any sequence $\{t_k\} \subset [0, 1)$ converging to 1, the sequence $\{\gamma(t_k)\}$ is special, respectively restricted.

Definition 2.3 ([1], [2]). Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary and let $z_0 \in D$. Let $p \in \partial D$ and $M > 1$. The *K-region* $K_{z_0}(p, M)$ of center τ , amplitude M and pole z_0 is

$$K_{z_0}(p, M) := \left\{ z \in D \mid \lim_{w \rightarrow p} [k_D(z, w) - k_D(z_0, w)] + k_D(z_0, z) < \log M \right\}.$$

Let $\{z_k\} \subset D$ be a sequence converging to $p \in \partial D$. It is known ([1, Lemma 2.7.12], [2]) that if $\{z_k\} \subset K_{z_0}(p, M)$ for some $M > 1$ then $\{z_k\}$ is restricted. While, if $\{z_k\}$ is special and restricted then it is eventually contained in a *K-region* $K_{z_0}(p, M)$ for some $M > 1$.

A holomorphic self-map f of D has K -limit q at $p \in \partial D$ —and we write $K\text{-}\lim_{z \rightarrow p} f(z) = q$ or, for short, $q = f(p)$ —if $\lim_{k \rightarrow \infty} f(w_k) = q$ for every sequence converging to p and such that $\{w_k\} \subset K_{z_0}(p, M)$ for some $M > 1$.

Definition 2.4. Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary and let $z_0 \in D$. Let $f: D \rightarrow D$ be holomorphic and let $p \in \partial D$. The *boundary dilation coefficient* $\alpha_p(f) \in (0, +\infty]$ of f at p is defined by

$$\frac{1}{2} \log \alpha_p(f) := \liminf_{w \rightarrow p} [k_D(z_0, w) - k_D(z_0, f(w))].$$

Since

$$(2.1) \quad \begin{aligned} k_D(z_0, w) - k_D(z_0, f(w)) &\geq k_D(f(z_0), f(w)) - k_D(z_0, f(w)) \\ &\geq -k_D(f(z_0), z_0) > -\infty, \end{aligned}$$

the boundary dilation coefficient is always strictly positive. Moreover, the boundary dilation coefficient does not depend on z_0 and can be computed using pluripotential theory as in the classical Julia's lemma for the unit disc (see [9]).

We state here the part of the Julia-Wolff-Carathéodory type theorem for strongly convex domains we need for our aims (see [1, Thm. 2.7.14], [2] for the first part, and [6, Prop. 3.4, Remark. 3.5] for (2)):

Theorem 2.5. *Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary and let $z_0 \in D$ and $p \in \partial D$. Let $f: D \rightarrow D$ be holomorphic and assume $\alpha_p(f) < +\infty$. Then $f(p) = K\text{-}\lim_{z \rightarrow p} f(z)$ exists and $f(p) \in \partial D$. Moreover,*

(1) *the holomorphic function*

$$D \ni z \mapsto \frac{1 - \tilde{\rho}_{f(p)}(f(z))}{1 - \tilde{\rho}_p(z)}$$

has limit $\alpha_p(f)$ along any special and restricted sequence $\{z_k\} \subset D$ which converges to p . In particular, the holomorphic self-map of $\mathbb{D}$ given by $\zeta \mapsto \tilde{\rho}_{f(p)}(f(\varphi_p(\zeta)))$ has a boundary regular fixed point at 1 and $\alpha_1(\tilde{\rho}_{f(p)} \circ f \circ \varphi_p) = \alpha_p(f)$.

(2) *The curve $[0, 1] \ni t \mapsto f(\varphi_p(t))$ converges to $f(p)$ and it is special and restricted.*

Boundary dilation coefficients satisfy the chain rule:

Lemma 2.6. *Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary. Let $f, g: D \rightarrow D$ be holomorphic and let $p \in \partial D$. If $\alpha_p(g \circ f) < +\infty$, then $\alpha_p(f) < +\infty$ and $\alpha_{f(p)}(g) < +\infty$, where $f(p) = K\text{-}\lim_{z \rightarrow p} f(z)$. Moreover,*

$$(2.2) \quad \alpha_p(g \circ f) = \alpha_{f(p)}(g) \cdot \alpha_p(f).$$

Conversely, if $\alpha_p(f) < +\infty$, then (2.2) holds.

Proof. Let $z_0 \in D$. Assume $\alpha_p(g \circ f) < +\infty$. Since

$$\begin{aligned} \alpha_p(g \circ f) &= \liminf_{w \rightarrow p} [k_D(z_0, w) - k_D(z_0, g(f(w)))] \\ &\geq \liminf_{w \rightarrow p} [k_D(z_0, w) - k_D(z_0, f(w))] \\ &\quad + \liminf_{w \rightarrow p} [k_D(z_0, f(w)) - k_D(z_0, g(f(w)))] \end{aligned}$$

and both terms in the right hand side are not $-\infty$ by (2.1), it follows that $\alpha_p(f) < +\infty$. By Theorem 2.5, $f(p)$ exists and $\alpha_{f(p)}(g) < +\infty$. Moreover, for $t \in [0, 1]$

$$(2.3) \quad \frac{1 - \tilde{\rho}_{g(f(p))}(g(f(\varphi_p(t))))}{1 - t} = \frac{1 - \tilde{\rho}_{g(f(p))}(g(f(\varphi_p(t))))}{1 - \tilde{\rho}_{f(p)}(f(\varphi_p(t)))} \frac{1 - \tilde{\rho}_{f(p)}(f(\varphi_p(t)))}{1 - t}.$$

Since $\tilde{\rho}_p(\varphi_p(t)) = t$, equation (2.2) follows by Theorem 2.5 letting $t \rightarrow 1$.

Conversely, assume $\alpha_p(f) < +\infty$. If $\alpha_{f(p)}(g) = +\infty$ then $\alpha_p(g \circ f) = +\infty$ for what we already proved. So we can assume $\alpha_{f(p)}(g) < +\infty$. By (2.3) and Theorem 2.5 the term on the left hand side has limit $\alpha_{f(p)}(g) \cdot \alpha_p(f)$ for $t \rightarrow 1$. By [9, Thm. 2.7] it follows that $\alpha_p(g \circ f) < +\infty$ and in fact (2.2) holds. $\square$

Definition 2.7. Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary. If $f: D \rightarrow D$ is holomorphic, we say that $p \in \partial D$ is a *contact point* for f if $K\text{-}\lim_{z \rightarrow p} f(z) = f(p)$ exists and $f(p) \in \partial D$. The contact point p is a *regular contact point* if $\alpha_f(p) < +\infty$.

In case $p = f(p)$, the point p is called a *boundary fixed point*. If it is also regular, it is called a *boundary regular fixed point*—or BRFP for short. For $A > 0$ we denote by

$$\text{BRFP}_A(f) := \{p \in \partial D \mid f(p) = p, \alpha_f(p) \leq A\}.$$

A point $p \in \text{BRFP}_A(f) \setminus \text{BRFP}_1(f)$ is called a *boundary repelling fixed point*.

A point $p \in \text{BRFP}_A(f)$ is called *isolated* if there exists a neighborhood U of p such that $\text{BRFP}_A(f) \cap U = \{p\}$.

In the unit disc, as a consequence of Cowen-Pommerenke's estimates [14] (see also [7, Thm. 2.2]), every boundary repelling fixed point is isolated. In higher dimension, this is no longer true (see [24, Example 6.3]).

The Denjoy-Wolff type theorem for strongly convex domains (see [1, Thm. 2.4.23], [9, Prop. 2.9] and [4]) can be stated as follows:

Theorem 2.8. *Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary and let $f: D \rightarrow D$ be holomorphic. Then either*

- (1) *there exists a complex geodesic $\varphi: \mathbb{D} \rightarrow D$ such that $f(\varphi(\zeta)) = \varphi(e^{i\theta}\zeta)$ for some $\theta \in \mathbb{R}$, and in particular $f(\varphi(0)) = \varphi(0)$ and $\alpha_f(p) = 1$ for all $p \in \partial\varphi(\mathbb{D})$, or*
- (2) *there exists $x \in D$ such that $f(x) = x$ and for every $p \in \partial D$ it holds $\alpha_f(p) > 1$ — and in such a case the sequence of iterates $\{f^{om}\}$ converges uniformly on compacta to the constant map $z \mapsto x$, or*
- (3) *$\text{BRFP}_1(f)$ contains a unique point $\tau \in \partial D$ such that the sequence of iterates $\{f^{om}\}$ converges uniformly on compacta to the constant map $z \mapsto \tau$ and $\alpha_f(p) > 1$ for all $p \in \partial D \setminus \{\tau\}$.*

A holomorphic self-map f of D is called *rotational elliptic* if it satisfies (1) of Theorem 2.8. It is called *strongly elliptic* if it satisfies (2) of Theorem 2.8. Finally, f is called *non-elliptic* if it satisfies (3) of Theorem 2.8.

In the non-elliptic case, the point τ is called the *Denjoy-Wolff point* of f , and f has no fixed points in D . In the strongly elliptic case, f has a unique fixed point in D .

Boundary regular stationary points, namely boundary regular fixed points with dilation coefficients = 1 are very special. The following lemma follows at once from Theorem 2.8 and [9, Prop. 2.9.(1)]:

Lemma 2.9. *Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary and let $f: D \rightarrow D$ be holomorphic. Let $p \in \partial D$ be such that $\alpha_f(p) = 1$. If p is isolated then f is non-elliptic and p is in fact the Denjoy-Wolff point of f .*

3. BACKWARD ITERATION SEQUENCES

In order to prove our main result, we shall use the so-called backward iteration sequences.

Definition 3.1. Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary and let $f: D \rightarrow D$ be holomorphic. A *backward iteration sequence* for f at $p \in \partial D$ is a sequence $\{w_k\} \subset D$ such that $f(w_{k+1}) = w_k$ for all $k \in \mathbb{N}$ and

$$\frac{1}{2} \log s(\{w_k\}) := \sup_{k \in \mathbb{N}} k_D(w_k, w_{k+1}) < +\infty.$$

The number $s(\{w_k\})$ is called the *hyperbolic step* of the sequence $\{w_k\}$.

Backward iteration sequences in the unit disc have been introduced in [7], exploiting results from [25], in order to study BRFP's for commuting mappings, and they have been thoroughly studied in [26]. Such results have been generalized to the ball in [24] and to strongly convex domains for non rotational elliptic maps in [3]. The following lemma is the content of [3, Lemma 2.2, Lemma 2.3]

Lemma 3.2. *Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary, and let $f: D \rightarrow D$ be a holomorphic map. Let $\{w_n\}$ be a backward iteration sequence converging toward the boundary of D . Then there exists a BRFP $p \in \partial D$ for f such that $w_n \rightarrow p$ and $\alpha_f(p) \leq s(\{w_k\})$.*

The following result is proved in [3, Thm. 0.1, Lemma 2.3, Thm. 3.3]:

Theorem 3.3. *Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary, and let $f: D \rightarrow D$ be a holomorphic map either strongly elliptic or non-elliptic.*

- (1) *If $\{w_k\}$ is a backward iteration sequence then $\{z_k\}$ converges to $p \in \text{BRFP}_A(f)$ for some $A \geq 1$ and $\alpha_f(p) \leq s(\{w_k\})$. In case $\alpha_f(p) = 1$ then f is non-elliptic and p is the Denjoy-Wolff point of f .*
- (2) *If $p \in \partial D$ is an isolated boundary repelling fixed point of f , then there exists a backward iteration sequence $\{w_k\}$ converging to p with hyperbolic step $s(\{w_k\}) \leq \alpha_f(p)$.*
- (3) *If $\{w_k\}$ is a backward iteration sequence which converges to p and $\alpha_f(p) > 1$, then there exists $M > 1$ and $k_0 \in \mathbb{N}$ such that $\{w_k\}_{k \geq k_0} \subset K_{z_0}(p, M)$.*

We examine now the rotational elliptic case. In order to state our result, we need some notations. Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary and let $f: D \rightarrow D$ be a holomorphic rotational elliptic map. By [1, Thm. 2.1.29] there exists a closed complex submanifold $M \subset D$ and a holomorphic map $\pi: D \rightarrow D$ such that $\pi(D) = M$, $\pi \circ \pi = \pi$, $\pi \circ f = f \circ \pi$, $f(M) = M$ and $f|_M$ is an automorphism of M . The manifold M is called the *limit manifold* of f since for all $z \in D \setminus M$ the limit set of $\{f^m(z)\}$ belongs to M . In particular, if $Z = \{z \in D \mid f(z) = z\}$, it follows that $\emptyset \neq Z \subseteq M$. The map π is called the *limit retraction* of f , and it can be obtained as limit of a sequence of iterates of f ; and moreover if $\pi': D \rightarrow D$ is another limit of a sequence of iterates of f such

that $\pi' \circ \pi' = \pi'$ then $\pi' = \pi$. In particular, all iterates of f have the same limit retraction and the same limit manifold.

Proposition 3.4. *Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary, and let $f: D \rightarrow D$ be a rotational elliptic holomorphic map and let M be the limit manifold of f .*

- (1) *A point $p \in \partial D$ is a regular contact point of f with $\alpha_f(p) = 1$ if and only if $p \in \overline{M} \cap \partial D$.*
- (2) *If $\{w_k\} \subset D$ is a backward iteration sequence and there exists $R > 0$ such that $\inf_{y \in M} \|w_n - y\| \geq R$ then $\{w_k\}$ converges to $p \in \text{BRFP}_A(f)$ for some $A > 1$ and $s(\{z_k\}) \geq \alpha_f(p) > 1$.*
- (3) *If $p \in \partial D$ is an isolated boundary repelling fixed point of f , then there exists a backward iteration sequence $\{w_k\}$ converging to p with hyperbolic step $s(\{w_k\}) \leq \alpha_f(p)$.*
- (4) *If $\{w_k\}$ is a backward iteration sequence which converges to p and $\alpha_f(p) > 1$, then there exists $M > 1$ and $k_0 \in \mathbb{N}$ such that $\{w_k\}_{k \geq k_0} \subset K_{z_0}(p, M)$.*

Proof. (1) Let $p \in \partial D \cap \overline{M}$. Let $z_0 \in M$ be such that $f(z_0) = z_0$, and let $\varphi_p: \mathbb{D} \rightarrow D$ be the complex geodesic such that $\varphi_p(0) = z_0$, $\varphi_p(1) = p$.

We claim that $\varphi_p(\mathbb{D}) \subset M$. Since $\pi \circ \pi = \pi$ and by the decreasing property of the Kobayashi distance, for all $z, w \in M$

$$k_M(z, w) = k_M(\pi(z), \pi(w)) \leq k_D(z, w) \leq k_M(z, w),$$

from which it follows that M is totally geodesic in D , i.e., $k_D|_M = k_M$. In particular, if $\eta: \mathbb{D} \rightarrow D$ is a complex geodesic such that $\eta(0) \in M$ and $\eta(r) \in M$ for some $r \in (0, 1)$, we have

$$k_D(\pi(\eta(0)), \pi(\eta(r))) = k_D(\eta(0), \eta(r)) = k_{\mathbb{D}}(0, r),$$

and hence $\pi \circ \eta: \mathbb{D} \rightarrow D$ is also a complex geodesic. By the uniqueness of complex geodesics up to pre-composition with automorphisms of $\mathbb{D}$, it follows that in fact $\pi \circ \eta(\zeta) = \eta(\zeta)$ for all $\zeta \in \mathbb{D}$. Thus, $\eta(\mathbb{D}) \subset M$ and η is a complex geodesic both for D and for M . Now, let $\{w_m\} \subset M$ be a sequence which converges to p and let $\varphi_m: \mathbb{D} \rightarrow D$ be the complex geodesic such that $\varphi_m(0) = z_0$ and $\varphi_m(r_m) = w_m$ for some $r_m \in (0, 1)$. For what we just proved, $\varphi_m(\mathbb{D}) \subset M$. Up to subsequences, we can assume that $\{\varphi_m\}$ converges uniformly on compacta to a holomorphic map $h: \mathbb{D} \rightarrow M$ such that $h(0) = z_0$. By Proposition 2.1, $\varphi_m \rightarrow h$ uniformly on $\overline{\mathbb{D}}$ and h is a complex geodesic for D (and for M). But, since $\varphi_m(r_m) \rightarrow p$, it follows that $h(1) = p$ and hence $h = \varphi_p$, which proves that $\varphi_p(\mathbb{D}) \subset M$.

Now, since $f: M \rightarrow M$ is an automorphism, and in particular an isometry for the Kobayashi distance, $f \circ \varphi_p: \mathbb{D} \rightarrow M$ is a complex geodesic in M , and hence in D . In particular, $\lim_{r \rightarrow 1} f(\varphi_p(r)) = q \in \partial D$ exists and $\varphi_q \equiv f \circ \varphi_p$. Therefore, $\tilde{\rho}_q(f(\varphi_p(\zeta))) = \zeta$ for all $\zeta \in \mathbb{D}$ and by Theorem 2.5.(1), $\alpha_f(p) = 1$.

Conversely, assume $p \in \partial D$ is a regular contact point with $\alpha_f(p) = 1$. Let $q = f(p)$ and $z_0 \in Z$. Let φ_p, φ_q be the complex geodesics such that $\varphi_p(0) = \varphi_q(0) = z_0$ and $\varphi_p(1) = p$, $\varphi_q(1) = q$. Then $g := \tilde{\rho}_q \circ f \circ \varphi_p$ is a holomorphic self-map of $\mathbb{D}$ such that $g(0) = 0$ and $g(1) = 1$, $\alpha_g(1) = \alpha_f(p) = 1$ by Theorem 2.5. As a consequence of the classical Julia's lemma [1, Corollary 1.2.10], $g(\zeta) = \zeta$ for all $\zeta \in \mathbb{D}$. Hence for all $\zeta_1, \zeta_2 \in \mathbb{D}$

$$k_{\mathbb{D}}(\zeta_1, \zeta_2) \leq k_D(f(\varphi_p(\zeta_1)), f(\varphi_p(\zeta_2))) \leq k_D(\varphi_p(\zeta_1), \varphi_p(\zeta_2)) = k_{\mathbb{D}}(\zeta_1, \zeta_2),$$

and thus $f \circ \varphi_p$ is a complex geodesic. The result follows then from:

Claim A: if $\varphi: \mathbb{D} \rightarrow D$ is a complex geodesic such that $\varphi(0) = z_0 \in Z$ and $f \circ \varphi: \mathbb{D} \rightarrow D$ is also a complex geodesic, then $\varphi(\mathbb{D}) \subset M$.

In order to prove Claim 1., recall that by [1, Thm. 2.1.21], the tangent space to D at z_0 admits a df_{z_0} -invariant splitting $T_{z_0}D = L_N \oplus L_U$ such that $T_{z_0}M = L_U$ and the spectrum of df_{z_0} in L_U is contained in $\partial\mathbb{D}$, while the spectrum of df_{z_0} in L_N is contained in $\mathbb{D}$. Note that $d\pi_{z_0}|_{L_U} = \text{id}$. Therefore if $\eta: \mathbb{D} \rightarrow D$ is a complex geodesic such that $\eta(0) = z_0$ and $\eta'(0) \in L_U$, then $\pi \circ \eta: \mathbb{D} \rightarrow D$ is a holomorphic map such that $\kappa_D(z_0; (\pi \circ \eta)'(0)) = \kappa_{\mathbb{D}}(0; 1)$, hence it is an infinitesimal isometry for the Kobayashi metric. Thus $\pi \circ \eta$ is a complex geodesic, and by the uniqueness of infinitesimal isometry, $\pi \circ \eta = \eta$. Hence $\eta(\mathbb{D}) \subset M$.

In the hypothesis of Claim A, $\kappa_D(z_0; \varphi'(0)) = \kappa_D(z_0; df_{z_0}(\varphi'(0)))$. Hence $\varphi'(0)$ belongs to L_U and therefore $\varphi(\mathbb{D}) \subset M$.

(2) *Step 1.* Let $z_0 \in Z$. We claim that

$$(3.1) \quad k_D(z, z_0) > k_D(f(z), z_0) \quad \forall z \in D \setminus M.$$

Clearly, $k_D(f(z), z_0) \leq k_D(z, z_0)$. Assume by contradiction that $k_D(f(z), z_0) = k_D(z, z_0)$ for some $z \in D \setminus M$. Let $\varphi: \mathbb{D} \rightarrow D$ be the complex geodesic such that $\varphi(0) = z_0$ and $\varphi(r) = z$ for some $r \in (0, 1)$. Then

$$k_D(f(\varphi(r)), \varphi(0)) = k_D(f(z), z_0) = k_D(z, z_0) = k_D(\varphi(r), \varphi(0)),$$

and therefore $f \circ \varphi$ is a complex geodesic in D . By Claim A, $\varphi(\mathbb{D}) \subset M$, thus $z \in M$, a contradiction.

Step 2. Let $z_0 \in Z$.

Claim B: For all $R_0 > 0$ there exists $0 < c = c(R_0) < 1$ such that for all $z \in D$ with $\inf_{y \in M} \|z - y\| \geq R_0$, it holds

$$(3.2) \quad k_D(f(z), z_0) - k_D(z, z_0) \leq \frac{1}{2} \log c < 0.$$

Assume (3.2) is not true. Then for every $c < 1$ there exists $z(c) \in D$ such that $\inf_{y \in M} \|z(c) - y\| \geq R_0$ and

$$(3.3) \quad k_D(f(z(c)), z_0) - k_D(z(c), z_0) > \frac{1}{2} \log c.$$

Let $x \in \overline{D}$ be a limit point of $\{z(1 - \frac{1}{n})\}$. Since $\inf_{y \in M} \|z(c) - y\| \geq R_0$, it follows that $x \notin \overline{M}$. If $x \in D$, it follows from (3.3) that $k_D(f(z(c)), z_0) \geq k_D(z(c), z_0)$, contradicting (3.1). Hence $x \in \partial D$ and

$$\liminf_{z \rightarrow x} [k_D(z, z_0) - k_D(f(z), z_0)] \leq 0.$$

But then x is a boundary regular contact point with $\alpha_f(x) \leq 1$, and by part (1), $x \in \overline{M}$, again a contradiction.

Step 3. Let $\{w_k\}$ be a backward iteration sequence satisfying the hypothesis of (2). Let $c = c(R)$ be given by Step 2. By induction on (3.2) we obtain that for all $k \in \mathbb{N}$,

$$e^{-2k_D(w_k, z_0)} \leq c^k e^{-2k_D(w_0, z_0)},$$

hence $k_D(w_k, z_0) \rightarrow \infty$ for $k \rightarrow \infty$. Therefore, $w_k \rightarrow \partial D$. The result follows then from Lemma 3.2.

(3) By (1), $p \notin \overline{M}$. Let $\epsilon > 0$ be such that $U := \{z \in \mathbb{C}^N \mid \|z - p\| < \epsilon\}$ has the property that $\overline{U} \cap \overline{M} = \emptyset$ and $U \cap \text{BRFP}_{\alpha_f(p)}(f) = \{p\}$. Arguing exactly

as in the proof of [3, Thm. 3.3] we can construct a backward iteration sequence $\{w_k\} \subset U$ with hyperbolic step $s(\{w_k\}) \leq \alpha_f(p)$. Since $\overline{U} \cap \overline{M} = \emptyset$, $\{w_k\}$ satisfies the hypothesis of (2), hence it converges to a boundary regular fixed point of f , say $q \in \partial D \cap U$, with $\alpha_f(q) \leq \alpha_f(p)$. But $U \cap \text{BRFP}_{\alpha_f(p)}(f) = \{p\}$ implies $q = p$ and we are done.

(4) Since $\{w_k\}$ converges to a boundary repelling fixed point $p \in \partial D$, and by (1), $p \notin \overline{M}$, hence $\{w_k\}$ satisfies the hypothesis of Claim B. Therefore, by (3.2)

$$\liminf_{k \rightarrow \infty} [k_D(z_0, z_{k+1}) - k_D(z_0, z_k)] \geq \frac{1}{2} \log \frac{1}{c} > 0.$$

Now the proof follows arguing exactly as in [3, Lemma 2.5]. □

4. SEMIGROUPS

Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary. A (continuous) semigroup (ϕ_t) of holomorphic self-maps of D is a continuous homomorphism from the semigroup $(\mathbb{R}^+, +)$ endowed with the Euclidean topology, to the semigroup (with respect of maps composition) of holomorphic self-maps of D endowed with the topology of uniform convergence on compacta (see, e.g., [27], [1], [9]).

If (ϕ_t) is a semigroup of holomorphic self-maps of D , we will denote by $\alpha_t(p) := \alpha_p(\phi_t)$ the dilation coefficient of ϕ_t at $p \in \partial D$.

Also, we denote by $\text{Fix}(\phi_t) := \{z \in D \mid \phi_t(z) = z \ \forall t \geq 0\}$.

The following result was proved in [1, Thm. 2.5.24, Prop. 2.5.26]; see also [5] and [8, Thm. A.1].

Theorem 4.1. *Let $D \subset \mathbb{C}^n$ be a bounded strongly convex domain with smooth boundary. Let (ϕ_t) be a semigroup of holomorphic self-maps of D . Then either*

- $\text{Fix}(\phi_t) \neq \emptyset$, or
- ϕ_t is non-elliptic for all $t > 0$ and there exists a unique $\tau \in \partial D$ such that τ is the Denjoy-Wolff point of ϕ_t .

The following result follows from [9, Corollary 4.8, Prop. 3.3]:

Proposition 4.2. *Let $D \subset \mathbb{C}^n$ be a bounded strongly convex domain with smooth boundary. Let (ϕ_t) be a semigroup of holomorphic self-maps of D .*

- (1) *Suppose there exists $t_0 > 0$ such that ϕ_{t_0} is strongly elliptic. Then ϕ_t is strongly elliptic for all $t \geq 0$.*
- (2) *If $p \in \partial D$ is a BRFP for ϕ_t for every $t \geq 0$, then there exists $\lambda \in (0, +\infty)$ such that $\alpha_t(p) = \lambda^t$.*

The previous results ensure that the following definition is well-posed:

Definition 4.3. Let $D \subset \mathbb{C}^n$ be a bounded strongly convex domain with smooth boundary. Let (ϕ_t) be a semigroup of holomorphic self-maps of D .

- (1) (ϕ_t) is *non-elliptic* if ϕ_1 is non-elliptic. In such a case, if $\tau \in \partial D$ is the Denjoy-Wolff point of ϕ_1 , we call τ the *Denjoy-Wolff point* of (ϕ_t) .
- (2) (ϕ_t) is *strongly elliptic* if ϕ_1 is strongly elliptic.
- (3) (ϕ_t) is *rotational elliptic* if ϕ_1 is rotational elliptic.

For rotational elliptic semigroups, (ϕ_t) , let M_t denote the limit manifold of ϕ_t for $t > 0$. We let

$$M(\phi_t) := M_1.$$

Lemma 4.4. *Let $D \subset \mathbb{C}^n$ be a bounded strongly convex domain with smooth boundary. Let (ϕ_t) be a rotational elliptic semigroup of holomorphic self-maps of D . Then $M_t = M(\phi_t)$ for every $t > 0$. In particular, $(\phi_t)|_{M(\phi_t)}$ is a group of automorphisms of $M(\phi_t)$.*

Proof. Since ϕ_1 is an iterate of $\phi_{1/q}$ for any $q \geq 1$, and the limit manifold of an iterate coincides with the limit manifold of the original map, we have $M_{1/q} = M_1$ for all $q \geq 1$. For the same reason we have $M_{p/q} = M_{1/q} = M_1$ for all $p/q \in \mathbb{Q}^+$. Since $\mathbb{Q}^+$ is dense in $\mathbb{R}^+$ it follows that $M_t = M_1$ for all $t > 0$. $\square$

5. REGULAR CONTACT POINTS FOR SEMIGROUPS

Contact points for semigroups of the unit disc were studied in [11] exploiting the existence of the so-called Königs' function, a tool which is not available in higher dimension. To replace it, we shall use complex geodesics.

Proposition 5.1. *Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary. Let (ϕ_t) be a semigroup of holomorphic self-maps of D . Let $t_0 > 0$ and let $p \in \partial D$ be a regular contact point for ϕ_{t_0} . Then there exists $T \in [t_0, +\infty]$ such that p is a regular contact point for ϕ_t for all $t \in [0, T)$. Moreover, the curve $[0, T) \ni r \mapsto \phi_r(p) \in \partial D$ is continuous.*

Proof. If p is the Denjoy-Wolff point of (ϕ_t) (when (ϕ_t) is non-elliptic) or it belongs to the closure of $\text{Fix}(\phi_t)$ (when (ϕ_t) is elliptic but not strongly elliptic) then p is a BRFP for all ϕ_t by Theorem 2.8 and the statement is true. Thus, we can suppose that p is neither the Denjoy-Wolff nor in the closure of $\text{Fix}(\phi_t)$.

Let $T := \sup t \in [0, +\infty)$ such that p is a regular contact point for ϕ_t . By assumption $T \geq t_0$.

Fix $\epsilon > 0$. If $T < +\infty$, let $T \geq t_1 > T - \epsilon$ be such that p is a regular contact point for ϕ_{t_1} . If $T = +\infty$, let $t_1 \geq 1/\epsilon$ be such that p is a regular contact point for ϕ_{t_1} . We will show that for all $t \in [0, t_1]$, p is a regular contact point for ϕ_t . Taking $\epsilon \rightarrow 0$, we will get the first statement.

To this aim, let $s, t > 0$ be such that $s + t = t_1$. Since $\phi_{t_1} = \phi_{t+s} = \phi_t \circ \phi_s$, it follows by Lemma 2.6 and Theorem 2.5 that p is a regular contact point for ϕ_s , that $\phi_s(p) \in \partial D$ is a regular contact point for ϕ_t and that

$$(5.1) \quad \alpha_t(\phi_s(p)) \cdot \alpha_s(p) = \alpha_{t+s}(p) = \alpha_{t_1}(p) < +\infty.$$

In particular, it follows that p is a regular contact point for ϕ_t for all $t \in [0, T)$.

Now, we consider the curve $[0, T) \ni r \mapsto \phi_r(p) \in \partial D$ and we prove that it is continuous. Fix $z_0 \in D$. Let $\varphi: \mathbb{D} \rightarrow D$ be the complex geodesic such that $\varphi(0) = z_0$ and $\varphi(1) = p$. Let $r_0 \in [0, T)$ be fixed. For $r \in [0, T)$ close to r_0 , let $\varphi_r: \mathbb{D} \rightarrow D$ denote the complex geodesic such that $\varphi_r(0) = z_0$ and $\varphi_r(1) = \phi_r(p)$ and let $\tilde{\varphi}_r: D \rightarrow \mathbb{D}$ be the left-inverse of φ_r . Let define

$$g_r(\zeta) := \tilde{\varphi}_r(\phi_r(\varphi(\zeta))) \quad \forall \zeta \in \mathbb{D}.$$

By construction, $g_r: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic and by Theorem 2.5.(1), 1 is a boundary regular fixed point for g_r with boundary dilation coefficient $\alpha_r(p)$. Thus, again by

the classical Julia-Wolff-Carathéodory theorem in $\mathbb{D}$, the derivative g'_r has non-tangential limit $\alpha_r(p)$ at 1.

Let fix $t_1 < T$ such that $t_1 > r_0$. From (5.1) we get that for all $r \leq t_1$ setting $t = t_1 - r$ it holds $\alpha_r(p) = \alpha_{t_1}(p)/\alpha_t(\phi_r(p))$. Now, if $\alpha_t(\phi_r(p)) \geq 1$ for all $r \in [0, t_1]$, it follows that $\alpha_r(p) \leq \alpha_{t_1}(p)$ for all $r \in [0, t_1]$. On the other hand, if $\alpha_t(\phi_{\tilde{r}}(p)) < 1$ for some $\tilde{r} \leq t_1$, by Theorem 2.8 it follows that $q = \phi_{\tilde{r}}(p)$ is the Denjoy-Wolff point of (ϕ_t) ; in particular, by Proposition 4.2, $\alpha_u(\phi_{\tilde{r}}(p)) = e^{\beta u}$ for all $u \geq 0$ and some $\beta < 0$ independent of $\tilde{r}$. It follows that $\alpha_{\tilde{r}}(p) \leq \alpha_{t_1}(p)e^{-\beta t_1}$, and thus we have proved that there exists $C > 0$ such that

$$\alpha_r(p) \leq C$$

for all $r \leq t_1$.

Let $M > 1$ and let

$$\overline{K} := \{\zeta \in \overline{\mathbb{D}} \mid |1 - \zeta| \leq M(1 - |\zeta|)\} \subset \mathbb{D} \cup \{1\}$$

be (the closure of) a Stolz angle in $\mathbb{D}$ with vertex 1 (see, e.g., [1, pag. 53]). Since $g'_r(1) = \alpha_r(p) \leq C$ for all $r \leq t_1$, it follows that $\{g'_r\}$ is equibounded in $\overline{K}$. Hence $\{g_r\}$ is equicontinuous on $\overline{K}$ (and it is clearly equibounded in $\overline{K}$ by 1). Applying Ascoli-Arzelà theorem, we find a subsequence $r_k \rightarrow r_0$ such that $\{g_{r_k}\}$ converges uniformly on $\overline{K}$ to some continuous function g . In particular, note that $g(1) = 1$.

Up to a subsequence, we can also assume that $\{\varphi_{r_k}\}$ converges uniformly on compacta of D to a holomorphic map $\eta: \mathbb{D} \rightarrow D$ such that $\eta(0) = z_0$. By Proposition 2.1, η is a complex geodesic and $\varphi_{r_k} \rightarrow \eta$ uniformly on $\overline{\mathbb{D}}$. Since $\varphi_{r_k}(1) = \phi_{r_k}(p)$ by construction, it follows that $\phi_{r_k}(p) \rightarrow \eta(1)$. Let $\tilde{\rho}: D \rightarrow \mathbb{D}$ be the left inverse of η . By Proposition 2.1, $\{\tilde{\rho}_{r_k}\}$ converges uniformly on $\overline{D}$ to $\tilde{\rho}$.

Since $\phi_{r_k} \rightarrow \phi_{r_0}$ uniformly on compacta of D , it follows that $g(\zeta) = \tilde{\rho}(\phi_{r_0}(\varphi(\zeta)))$ for all $\zeta \in K$. Taking the radial limit at 1, we obtain

$$1 = \lim_{(0,1) \ni s \rightarrow 1} g(s) = \lim_{(0,1) \ni s \rightarrow 1} \tilde{\rho}(\phi_{r_0}(\varphi(s))) = \tilde{\rho}(\phi_{r_0}(p)).$$

Since the only point in the fiber of $\tilde{\rho}$ over 1 is the point $\eta(1)$, it follows that $\eta(1) = \phi_{r_0}(p)$, hence $\phi_{r_k}(p) \rightarrow \phi_{r_0}(p)$. Repeating the argument for any subsequence, we obtain the result. $\square$

6. COMMON BRFP'S FOR SEMIGROUPS

Theorem 6.1. *Let $D \subset \mathbb{C}^N$ be a bounded strongly convex domain with smooth boundary. Let (ϕ_t) be a semigroup of holomorphic self-maps of D . Suppose $p \in \partial D$ is an isolated boundary repelling fixed point for ϕ_{t_0} for some $t_0 > 0$. Then p is a boundary repelling fixed point for ϕ_t for all $t > 0$.*

Proof. By Theorem 3.3 or Proposition 3.4 there exists a backward iteration sequence $\{w_n\}$ for ϕ_{t_0} such that $k_D(w_n, w_{n+1}) \leq \frac{1}{2} \log \alpha_{t_0}(p)$ and $\{w_n\}$ converges to p inside a K -region. For $t \in [0, t_0]$, define $z_n^t := \phi_t(w_n)$. Then

$$\phi_{t_0}(z_n^t) = \phi_t(\phi_{t_0}(w_n)) = \phi_t(w_{n+1}) = z_{n+1}^t,$$

and

$$(6.1) \quad k_D(z_n^t, z_{n+1}^t) = k_D(\phi_t(w_n), \phi_t(w_{n+1})) \leq k_D(w_n, w_{n+1}) \leq \frac{1}{2} \log \alpha_{t_0}(p).$$

Hence $\{z_n^t\}$ is a backward iteration sequence for ϕ_{t_0} . Moreover, by Proposition 5.1, p is a regular contact point for ϕ_t all $t \in [0, t_0]$. Since $\{w_n\}$ converges to p inside a K -region, by Theorem 2.5

$$q_t := \lim_{n \rightarrow \infty} z_n^t = \lim_{n \rightarrow \infty} \phi_t(w_n) = \phi_t(p) \in \partial D.$$

Therefore, by Lemma 3.2, q_t is a BRFP for ϕ_{t_0} with dilation coefficient $\alpha_{t_0}(q_t) \leq \alpha_{t_0}(p)$.

Hence, by Proposition 5.1, the curve $[0, t_0] \ni t \mapsto q_t$ is a continuous curve made of BRFP's of ϕ_{t_0} and $\alpha_{t_0}(q_t) \leq \alpha_{t_0}(p)$. Since p is isolated, the only possibility is $q_t = p$ for all $t \in [0, t_0]$.

Now, let $t > 0$. Then $t = mt_0 + s$ for some $m \in \mathbb{N}$ and $s \in [0, t_0]$. Hence

$$\phi_t(p) = \phi_{mt_0+s}(p) = \phi_{mt_0}(\phi_s(p)) = \phi_{t_0}^m(p) = p.$$

Moreover, by Lemma 2.6, $\alpha_t(p) = \alpha_{t_0}(p)^m \cdot \alpha_s(p) < +\infty$, which implies that p is a common BRFP for (ϕ_t) . $\square$

Now we can prove Theorem 1.1:

Proof of Theorem 1.1. By Theorem 4.1 and Lemma 2.9, if $\alpha_f(p) \leq 1$ then p is the common Denjoy-Wolff point of (ϕ_t) . If $\alpha_f(p) > 1$ the result follows from Theorem 6.1. $\square$

For rotational elliptic semigroups there might exist boundary regular (in fact stationary) non isolated fixed points which are not fixed for all the elements of the semigroup:

Example 6.2. Let $\phi_t(z, w) = (e^{2\pi i t} z, w)$. Then (ϕ_t) is a rotational elliptic semigroup of $\mathbb{B}^2$. The points $\partial \mathbb{D} \times \{0\}$ are BRFP's for ϕ_1 (with boundary dilation coefficient 1) but not for ϕ_t with $t \not\equiv 0 \pmod{1}$.

REFERENCES

- [1] M. Abate, *Iteration theory of holomorphic maps on taut manifolds*, Mediterranean Press, Rende, 1989.
- [2] M. Abate *The Lindelöf principle and the angular derivative in strongly convex domains*. J. Anal. Math, 54 (1990), 189-228.
- [3] M. Abate, J. Raissy, *Backward iteration in strongly convex domains*. Adv. Math. 228 (2011), 2837-2854.
- [4] M. Abate, J. Raissy, *Wolff-Denjoy theorems in non-smooth convex domains*. To appear in Ann. Mat. Pura Appl., arXiv: 1211.2454! (2013).
- [5] L. Aizenberg and D. Shoikhet, *Boundary behavior of semigroups of holomorphic mappings on the unit ball in $\mathbb{C}^n$* , Complex Variables Theory Appl. **47** (2002), 109-121.
- [6] F. Bracci, *Commuting holomorphic maps in strongly convex domains*, Annali della Scuola Normale Sup. di Pisa, Cl. di Sc. (4) 27 (1998), 1, 131-144.
- [7] F. Bracci, *Fixed points of commuting holomorphic mappings other than the Wolff point*. Trans. Amer. Math. Soc., 355, 6, (2003), 2569-2584.
- [8] F. Bracci, M. D. Contreras, S. Díaz-Madrigal, *Classification of semigroups of linear fractional maps in the unit ball*. Adv. Math., 208, (2007), 318-350.
- [9] F. Bracci, M. D. Contreras, S. Díaz-Madrigal, *Pluripotential theory, semigroups and boundary behavior of infinitesimal generators in strongly convex domains*. J. Eur. Math. Soc. 12 (2010), 23-53.
- [10] F. Bracci, G. Patrizio, S. Trapani, *The pluricomplex Poisson kernel for strongly convex domains*. Trans. Amer. Math. Soc., 361, 2, (2009), 979-1005.
- [11] F. Bracci, P. Gumenyuk, *Contact points and fractional singularities for semigroups of holomorphic self-maps in the unit disc*. arXiv:1309.2813

- [12] M.D. Contreras, S. Díaz-Madrigal, Ch. Pommerenke, *Fixed points and boundary behaviour of the Koenigs function*. Ann. Acad. Sci. Fenn. Math. 29, (2004), 471—488.
- [13] M. D. Contreras, S. Díaz-Madrigal, Ch. Pommerenke, *On boundary critical points for semigroups of analytic functions*. Math. Scand. 98 (2006), no. 1, 125—142.
- [14] C.C. Cowen and Ch. Pommerenke, *Inequalities for the angular derivative of an analytic function in the unit disk*. J. London Math. Soc. (2), 26 (1982), 271- 289.
- [15] M. Elin, S. Reich, and D. Shoikhet, *Asymptotic behavior of semi- groups of ρ -nonexpansive and holomorphic mappings on the Hilbert ball*, Ann. Math. Pura Appl. (4) ,181, (2002), 501–526.
- [16] M. Elin, D. Shoikhet, *Semigroups of holomorphic mappings with boundary fixed points and spirallike mappings*. Geometric function theory in several complex variables, 82–117, World Sci. Publ., River Edge, NJ, 2004.
- [17] M. Elin, S. Reich, and D. Shoikhet, *A Julia-Caratheodory theorem for hyperbolically monotone mappings in the Hilbert ball*, Israel J. Math. 164 (2008), 397411.
- [18] M. Elin and D. Shoikhet, *Dynamic extension of the Julia-Wolff Carathéodory theorem*, Dynam. Systems Appl. **10** (2001), 421–438.
- [19] M. Elin and D. Shoikhet, *Semigroups of holomorphic mappings with boundary fixed points and spirallike mappings*, Geometric function theory in several complex variables, 82–117, World Sci. Publishing, River Edge, NJ, 2004.
- [20] S. Kobayashi, *Hyperbolic complex spaces*. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 318. Springer-Verlag, Berlin, 1998. xiv+471 pp. ISBN: 3-540-63534-3
- [21] L. Lempert, *La métrique de Kobayashi et la représentation des domaines sur la boule*. Bull. Soc. Math. Fr. 109 (1981), 427-474.
- [22] L. Lempert, *Holomorphic retracts and intrinsic metrics in convex domains*. Analysis Math. 8 (1982), 257-261.
- [23] L. Lempert, *Intrinsic distances and holomorphic retracts*. Complex Analysis and Applications'81, Sofia, (1984), 341-364.
- [24] O. Ostapyuk, *Backward iteration in the unit ball*. Illinois J. Math. 55 (2011), no. 4, 15691602 (2013).
- [25] P. Poggi Corradini, *Canonical conjugation at fixed points other than the Denjoy-Wolff point*. Ann. Acad. Sci. Fenn. Math. 25 (2000), 2, 487-499.
- [26] P. Poggi Corradini, *Backward sequences with bounded hyperbolic steps for analytic self-maps of the disk*. Revista Matematica Iberoamericana, 19 (2003), no. 3, 943970.
- [27] S. Reich, D. Shoikhet, *Nonlinear semigroups, fixed points, and geometry of domains in Banach spaces*. Imperial College Press, London, 2005. xvi+354 pp. ISBN: 1-86094-575-9
- [28] A. G. Siskakis, *Semigroups of Composition Operators and the Cesàro Operator on $H^p(D)$* . Ph.D. Thesis, University of Illinois, 1985.

M. ABATE: DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, L.GO PONTECORVO 5, 56127 PISA, ITALY

E-mail address: `abate@dm.unipi.it`

F. BRACCI: DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA “TOR VERGATA”, VIA DELLA RICERCA SCIENTIFICA 1, 00133, ROMA, ITALY.

E-mail address: `fbracci@mat.uniroma2.it`